

Flexhaler® Fact Sheet

Pulmicort® is an oral corticosteroid medication intended for the maintenance and treatment of asthma in adults and children 6 years of age and older. Budesonide is not a bronchodilator and is not intended for the treatment of acute asthma exacerbation. The flexhaler® contains an anti-inflammatory medicine (budesonide) which reduces airway swelling, inflammation, and irritation.

Generic Name: Budesonide

Brand Name: Pulmicort®

Dosage Administration and Strength

90 Pulmicort® Flexhaler®: delivers 90 mcg budesonide inhalation powder

• 180 Pulmicort® Flexhaler®: delivers 180 mcg budesonide inhalation powder

Indications

Adults and children 6 years of age or older with asthma

Contraindications

 Flexhaler® should not be used for sudden symptoms of shortness of breath or acute episodes of asthma

Side Effects

- Inflammation of the nose and throat
- Nasal congestion and nasal allergies
- Viral upper respiratory tract infection
- Thrush in the mouth and throat
- Rare instance of increased intraocular pressure, cataracts and glaucoma

Cleaning Instructions

- Do not immerse in water, keep the inhaler dry at all times
- To clean, wipe inhaler with a dry towel or tissue
- Store inhaler in a dry place and away from moisture at all times

References

- 1. American Lung Association." *Asthma Medicines Chart*. Revised Jan. 2009 Retrieved from www.lung.org.
- 2. Pulmicort® Flexhaler® [prescribing Information]. Wilmington, DE. AstraZeneca; 2014.



Patient Name:
Patient ID:
Evaluator Initials:
Date:/

PULMICORT® FLEXHALER® INHALATION TECHNIQUE CHECKLIST <u>Trained assessor to read to participant</u>: Please show me exactly how you use your Pulmicort® Flexhaler® at home.

	Pre		Post 1		Post 2		Post 3		
	Wrong	Correct	Wrong	Correct	Wrong	Correct	Wrong	Correct	
Twists and removes cover			\square_0		\square_0		\square_0		
2. Holds the inhaler upright with mouthpiece facing up					\square_0				
3. Twists the grip to the right then left until it clicks									
4. Breathes out fully					\Box_0		\Box_0		
5. When breathing out fully, does so away from inhaler			\Box_0		\Box_0		\Box_0		
Puts inhaler into mouth, closes lips around mouthpiece									
7. Breathes in deeply and forcefully									
Holds breath for ten seconds (with or without inhaler in mouth)									
9. Removes inhaler before breathing normally					\square_0		\square_0		
10. Places cover back on inhaler									
11. Twists to close					\square_0		\square_0		
12. Rinses mouth									
Participant's Score:		/12		/12		/12		/12	
	[] N/A		[] N/A		[] N/A		[] N/A		

Score the participant's performance by assigning 1 point to each correct step and adding across the 12 steps. If score is less than 12, demonstrate to the participant appropriate technique and repeat inhaler assessment.

Comments:			